LIST OF OUR POSTERS




Clinical profile of community-acquired pneumonia (CAP) and vaccination status among children <5 years in public & private sector: A retrospective review of inpatients records.

ECCMID 26th Amsterdam

Cost Effectiveness of Pegylated Interferon α -2b in Previously Untreated Chronic HCV Patients in Mizoram, India

Annual Conference of the Indian National Association



The Control Control Scott Scot

Direct medical cost associated with the diagnosis and treatment of patients with chronic hepatitis-b in three large metropolitan cities in India

ISPOR 18th Annual European

Relationship of pen needle reimbursement (PNR) to health resource utilization (HRU) and costs in Chinese diabetes mellitus (DM) patients

ВD

Endocrinology, Sautheast University Zhongda Hospital, Narijing, China; [†] Endocrinology, Peking University Peop 8D and Company, Beijing, China		
Abstract		
AA affects an estimated 314 million people in China with another 497 million classified as pre-DM. LH is an addressable complication	+ Patients were mean \$9.5 (\$D\$1)	Siya sidand were t
mong insulin injecting DM patients that may affect insulin absorption and thereby related costs. This study realisates the prevalence of UK	Table 1. Overall Damographic Character	eletics
and it's relationship to insulin consumption in China. An observational study was conducted among 403 (AM patients from 4 cities (Beiling,	Diamounting	Statistics
sarying, Chongging, Zhengshou). Participants were continuously injecting injulin for a 1 year, 18-80 yrs old with a 8MI a 38.5 kg/lin ² .	Age (sear)	-
temperaphics, medical history direct and indirect costs and disease management data were collected via patient survey. Astroped to a	Mean DCI	98601.9
theirs) exam and HSAS; tests. Patients were meanSSS SHASTS I un old, and took insulin SASASI un, averaging 33.0/18.6/10/fes	Conde	
Prevalence of UPI was \$2.9%, PBA2c = 8.2(1.8)% in those with, and 7.3(1.5)% in those without UPI a=0.00(1). Patients with UPI used an average	Main	201 (40.3.5)
f [1] U/Say more in comparison with non-UH injectors [38,1 U/ss 27.1 U, av0.000] and were more likely to have HBA1; a 7% (58 % vs. 42%.	Semale	202 200 1 762
in C022). Daily injulin cost (RM8.8.2 yr.5.8, px2.001) and 80% rearmalized daily injulin cost (RM8.0.31 yr.0.24, px0.001) were higher in those	Acceptancement status of PN	
institute, start materials protest a visit protest for our force of where the average results note is RMM 215 per it. The result force of depression of the control of exercise of exercise of the control of exercise o	Tes	141205430
ens United are approximately 6.4 minor equal right 55 or Units, where the average making pice 0.6 minor (13 per 9 minor occurs equal realist see associated with UR is estimated to be approximately EMB 1.892,993,600 (56.00 million) per year. UR in insulin taking patients	No	259 844 93
ersents a significant, avaidable consents burden in Chine due to excess insulin assumption in the setting of worse physicial control money those affectual insplementing techniques to avoid or limit (II development could result in decreased routin expenditure, lower routing consumption, and may improve physicial control.	LH was provident in \$2.9% of the without UR 7.7% (SD±1.5)(p=0.0) Patients with LH used on an aver have MBAS levels a 7% (SBS vs.	035 (Table 2) rage 11 U of insulin /
Introduction	Table 2. Ownell Clinical Charac	Harristics by LH Charan
Disdutes excline ICM affects as estimated 116 million secretaris China with another 817 million also find as businesse eliabetes."	Parameter	
As more patients gain access to and inject insulin, UR is bound to become more prevaient. Previously reported prevaience of UR has been found to be as high as 64.85% in insulin injecting patients."	Type of dishetes (x=800) Type 1 Type 5	
Intent hand the learning as d.d. Cut misstering registeries. Unitered indicate the Cut for appreciating flow muscular shouthers and lead to problems with give area (corteq): United associated with improper improtor inchroque, total daily stope of impute, number of impostume a day and revue of needles. United associated with improper improtor inchroque, total daily stope of impute, number of impostume a day and revue of needles. United associated earth. United associated with improper improtor inchroque, total daily stope of impute, number of impostume a day and revue of needless. United associated earth. United associated earth improved impostume and impostume and impostume and impostume a day and revue of needless and impostume a	Type II Duration of time with DA/ Mount (I/O) Chance control statut < 70. rishit > 70. Frequency of Ingrafication	